

Treatment Start DATE:	Baseline	2 Week DATE:	4 Week DATE:	8 Week DATE:	12 Week DATE:	16 Week DATE:	20 Week DATE:	24 Week DATE:	28 Week* DATE:	32 Week* DATE:	36 Week* DATE:	40 Week* DATE:	44 Week* DATE:	48 Week* DATE:	12 week post-tx DATE:
<input type="text"/>		<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
Lab date															
WBC															
ANC															
Hgb															
Platelets															
Cr															
AST															
ALT															
HCV RNA QUANT															
TSH															
Assess depression															
Pregnancy Counseling															
Assess side effects															

TELAPREVIRPRESCRIBING RECOMMENDATIONS** (Abbreviations: TEL: telaprevir; PEG: pegylated interferon; RBV: ribavirin)

**If on treatment*

***Nonformulary*

PATIENT TYPE	WEEK 4 VIRAL LOAD	WEEK 12 VIRAL LOAD	TREATMENT DURATION	TOTAL TREATMENT DURATION
Treatment naive or Treatment experienced: prior relapsers	Undetectable	Undetectable	12 weeks TEL/PEG/RBV, then additional 12 weeks PEG/RBV	24 weeks
	≤1000 IU/ml	≤1000 IU/ml	12 weeks TEL/PEG/RBV, then additional 36 weeks PEG/RBV	48 weeks
Treatment naive with cirrhosis; OR Treatment naive or Treatment experienced HIV positive	Undetectable	Undetectable	12 weeks TEL/PEG/RBV, then additional 36 weeks PEG/RBV	48 weeks
Treatment experienced: prior partial responders and null responders	≤1000 IU/ml	≤1000 IU/ml	12 weeks TEL/PEG/RBV, then additional 36 weeks PEG/RBV	48 weeks

TELAPREVIR FUTILITY RULES

HCV VIRAL LOAD	ACTION
WEEK 4>1000 IU/ml	DISCONTINUE TREATMENT
WEEK 12>1000 IU/ml	DISCONTINUE TREATMENT
WEEK 24 any detectable viral load	DISCONTINUE TREATMENT

CDCR #:	
Last Name:	
First Name:	MI:
DOB:	